

Status of Reference Event Collection

István Bondár, Bob Engdahl, Hans
Israelsson, Xiaoping Yang, Rami Hofstetter,
Hafidh Ghalib, Indra Gupta, Bob Wagner,
Victor Kirichenko, Keith McLaughlin


DTRA Program Review
CMR, March 21-22, 2002

Overview

- Objectives
- Consortium effort to collect reference events
- Methodology
- Reference Event List 2.0
- Event clusters
- Summary

Objectives

- Compile a list of GT0-10 reference events in the Consortium's region of interest with emphasis on GT0-5 events
- Validate, document and QC each event
- Reference events are employed to validate
 - models
 - corrections and errors.



Consortium effort


- Nearly half of the Consortium work (and budget) is devoted to reference event collection and validation with the recognition that models and SSSCs cannot be validated without a suitable set of ground truth events
- Contacts have been established in the region (Multimax, GII, CUB, SAIC, WSC)
- Local bulletins that may not be publicly available have been acquired (Multimax, WSC, CUB, SAIC, Harvard)
- Phases are being picked from waveforms from non-reporting future IMS stations (GII, Multimax)
- Validation and documentation of reference events is an ongoing effort (CUB, SAIC)

Major contributions

<http://g2calibration.cmr.gov/calibration/acknowledgement.html>

<http://g2calibration.cmr.gov/calibration/refref.html>

- DOE LLNL delivery
- Local bulletins from Spain, Morocco, Kuwait, Saudi Arabia, Northern Caucasus, Turkey, Greece, Cyprus, Austria, Hungary, Romania, Egypt
- Shot locations of EGT, EuroBridge, Celebration2000, Vrancea99 and Saudi refraction profiles
- NNCKR list of Balapan nuclear explosions, PNE readings collected at Harvard, refined locations of Lop Nor explosions
- Reference events for clusters from Iran, China, India, Turkey, Egypt; potential clusters from Kuwait, Ghana, Djibouti
- Mid-ocean ridge and transform fault events
- ISC, NEIC, EMSC, REB, EHB searches
- CMR databases (Explo, REDB, Gamma, GT)




Methodology

- Identify candidate reference events
 - Bulletin search using GT5 selection criteria
 - Use other independent information (e.g. bathymetry, space imagery, published papers, mine location, personal communications, etc.)
- Collect and merge arrivals from various sources
 - GT selection criteria may not be met without merging bulletins
- Validate candidate reference events
 - Cluster analysis (HDC, JHD)
 - Relocation using only local stations
- QC and vet arrivals
 - Conflicting station codes
 - Possible phase misassociations

GT5 selection criteria (Bondár et al.)


Relocation of two GT0 events with 10 randomly selected stations within 250km; 10,000 realizations

- No universal selection criteria exists
- Criteria that minimize the number of outliers and maximize the number of events with 5 km location accuracy at the 90% and 95% confidence level




Secondary azimuthal gap

- The largest azimuthal gap filled by a single station
- Tends to equalize relative station importances



3/27/02





DTRA program review, 21-22 March, 2002

GT5 selection criteria



At the 95% confidence level

- At least 10 stations within 250 km with azimuthal gap ≤ 110 degrees and with secondary azimuthal gap ≤ 160 degrees
- At least one station within 30 km from the epicenter
- Event is recorded beyond 250 km




At the 90% confidence level

- At least 10 stations within 250 km with azimuthal gap ≤ 110 degrees
- At least one station within 30 km from the epicenter
- Event is recorded beyond 250 km




GT15 selection criteria (Engdahl et al.)

Relocation of 1,800 GT0-5 earthquakes and explosions using stations at all distances and later phases. Preliminary results.



Percent of mislocation of 15 km or less



Validation of reference events


- Each reference event is documented with metadata
- Cluster analysis (CUB, SAIC)
 - candidate events are validated if multiple event location results are consistent with reference event information
 - Events from cluster are promoted to GT5 if the major axis of their error ellipse is less than 5.5 km
 - HDC and JHD techniques were cross-validated to ensure they give similar results
- Relocation using stations within 300 km if clusters cannot be formed (Multimax)
- Oracle database is sponsored by CMR

Sources of arrival data

- Global bulletins: ISC, EHB, PDE, REB, REDB
- Local bulletins collected by
 - Multimax (Kuwait, Morocco, Saudi Arabia, Spain),
 - GII (Turkey, Jordan, Cyprus, Greece)
 - WSC in cooperation with GSRAS (Azerbaijan, Dagestan, Georgia)
 - SAIC (Austria, Hungary, Romania, Slovenia)
- Arrivals picked from stations at future IMS sites if they are not reported to ISC (GII, Multimax)
- Arrivals are merged and vetted
 - Conflicting station codes are corrected
 - Duplicate arrivals are deleted
 - Possible phase misassociations (Pg/Pn/P, Sg/Sn/S) are investigated

Reference Event List 2.0 March, 2002

1870 GT0-10 events (REL1.0: 837 events)



3/27/02

DTRA program review, 21-22 March, 2002


12

Reference Event List 2.0 March, 2002


64,258 regional rays (Pg, Pn, Sn, Lg)

126,360 teleseismic (P, S) rays

Limited coverage in Africa



Distribution of phases



Reference Event List 2.0

- The Reference Event List is publicly available
- Submitted to the IASPEI Working Group on Reference Events

<http://g2calibration.cmr.gov/calibration/data.html>

<http://g2calibration.cmr.gov/calibration/refsel.html>

Browse Reference Event List

	Minimum	Maximum	Include: <input checked="" type="checkbox"/> Metadata <input type="checkbox"/> Phases
date	<input type="text" value="1994/01/01"/>	<input type="text" value="1995/01/01"/>	
latitude	<input type="text" value="0"/>	<input type="text" value="20"/>	
longitude	<input type="text" value="40"/>	<input type="text" value="60"/>	
depth	<input type="text"/>	<input type="text"/>	
<div>Submit Query Reset</div>			

Reference Event List 2.0

Search results

For events between 1994/01/01 and 1995/01/01 in the coordinate rectangle (0.000, 40.000) - (20.000, 60.000)

Mid-Indian ridge	GT-level: 10	Author: PAN
Pan, J., M. Antolik and A. Dziewonski, Locations of mid-oceanic earthquakes constrained by sea-floor bathymetry, EOS Trans. AGU, 81, F868, 2000.		

DATA_TYPE ORIGIN IMS1.0

Reference Event List of the Group2 Consortium, generated 2002/03/14 21:01:37

Date	Time	Err	RMS	Latitude	Longitude	S _{max}	S _{min}	Az	Depth	Err	Ndef	Nsta	Gap	ndist	Mdist	Qual	Author	OrigID
1994/03/19	10:43:32.10		4.15	7.4400	58.4300				10.0		132	104	103	17.31	127.50	m i uk	GROUP2	20818554

Magnitude Err Nsta Author OrigID

(GT10- 10 km accuracy)

Gulf of Aden cluster	GT-level: 10	Author: ENGDAHL_HDC
Engdahl, E.R. and E.A. Bergman, Validation and generation of reference events by cluster analysis , 23rd Seismic Research Review, Vol I, 205-214, October 2-5, 2001.		

DATA_TYPE ORIGIN IMS1.0

Reference Event List of the Group2 Consortium, generated 2002/03/14 21:01:39

Date	Time	Err	RMS	Latitude	Longitude	S _{max}	S _{min}	Az	Depth	Err	Ndef	Nsta	Gap	ndist	Mdist	Qual	Author	OrigID
1994/10/01	14:04:22.15	0.15	1.23	13.1500	50.3350	2.1	5.4	280	10.0f	0.0	145	145	110	9.09	91.85	m i uk	GROUP2	21450306

Magnitude Err Nsta Author OrigID

mb	5.0		GROUP2	21450306
Ms	4.4		GROUP2	21450306

(GT10- 10 km accuracy)



3/27/02

DTRA program review, 21-22 March, 2002

15


Event cluster database

- Empirical path corrections produced by cluster analysis are used to validate SSSCs and model errors
- 62 event clusters, 4,004 regional and 7,691 teleseismic rays




Event cluster database

Path corrections vs distance




Variance of path corrections vs distance



Event cluster database

- Pairs of clusters: correlation of path corrections between close clusters at a station; might be biased by origin time errors
- Pairs of stations: correlation of path corrections between stations at a cluster
- Empirical path corrections provide estimate for typical correlation length: ~100 km



Summary

- Reference event collection, validation and documentation is a major Consortium effort
- Reference event collection requires extensive international cooperation
- Reference event selection criteria are established
- More than 1,800 GT0-10 events in current Reference Event List
- Reference Event List releases are publicly available at <http://g2calibration.cmr.gov>
- Event cluster database with empirical path corrections is created